





















Guerin-Danon, C., Chabanet, C., Pedone, C., Popot, F., Vaisside, P., Bouley, C., Szylit, O. & Andrieux, C. (1998). Milk fermented with yogurt cultures and *Lactobacillus casei* compared with yogurt and gelled milk: influence on intestinal microflora in healthy infants. *American Journal of Clinical Nutrition*, 67, 111-117.

Isolauri, E., Sutas, Y., Kankaapaa, P., Arvilommi, H. & Salminen, S. (2001). Probiotics: Effects on immunity. *American Journal of Clinical Nutrition*, 73, 444S-450S.

Kailasapathy, K., and Rybka, S. (1997). *Lactobacillus acidophilus* and *Bifidobacterium* ssp.-Their therapeutic potential and survival in yoghurt. *Australian Journal of Dairy Technology*, 52, 28-35

Liong, M. & Shah, N. P. (2005). Acid and bile tolerance and cholesterol removal ability of lactobacilli strains. *Journal of Dairy Science*, 88, 55-66.

Lourens-Hattingh, A. & Viljoen, B. C. (2001). Yoghurt as probiotic carrier food. Review article. *International Dairy Journal*, 11, 1-17

Mattila-Sandholm, T., Myllarinen, P., Crittenden, R., Mogensen, G., Fonden, P. & Saarela, M. (2002). Technological challenge for future probiotic foods. *International Dairy Journal*, 12, 173-182.

Oliveira, M. N., Sodini, I., Rezzonico, F. & Corrieu, G. (2001). Effect of milk supplementation and culture composition on acidification, textural properties and microbiological stability of fermented milks containing probiotic bacteria. *International Dairy Journal*, 11, 935-942.

Paseephol, T. (2008). Characterisation of prebiotic compounds from plant sources and food industry wastes. Inulin from Jerusalem artichoke and D-fructose from milk concentration permeate, 1-21.

Paftik, I. (2002). Lactic acid bacteria and cheese: Metabolic perspective. *British Journal of Nutrition*, 88 (Suppl.), S89-S94.

Rasooli, M., Parekh, T., Dave, N., Patel, V. & Subhash, R. (2008). Evaluation of various physico-chemical Properties of Hibiscus saffdarika and *L. casei* incorporated probiotic yogurt. *Pakistan Journal of Biological Sciences*, 11, (17), 2101-2108.

Saide, J. A. O. & Gilliland, S. E. (2005). Antioxidative activity of Lactobacilli measured

by oxygen radical absorbance capacity. *Journal of Dairy Science*, 88, (4), 1352-1357.

Shah, N. P. (2007). Functional cultures and health benefits. *International Dairy Journal*, 17, 1262-1277

Shah, N. P. (2000). Probiotic bacteria: Selective Enumeration and Survival in Dairy Foods. *Journal of Dairy Science*, 83, 894-907.

Shah, N. P. & Lankaputhra, W. E. V. (1997). Improving viability of *Lactobacillus acidophilus* and *Bifidobacterium bifidum* ssp. In yoghurt. *International Dairy Journal*, 7, 343-356.

Tharmaraj, N. & Shah, N. P. (2002). Selective Enumeration of *Lactobacillus delbrueckii* ssp. *Bulgarius*, *Streptococcus thermophilus*, *Lactobacillus acidophilus*, *Bifidobacteria*, *Lactobacillus casei*, *Lactobacillus lactis*, and *Propionibacteria*. *Journal of Dairy Science*, 86, 2288-2296.

Unal, B., Metin, S. & Iski, N. D. (2003). Use of response surface methodology to describe the combined effect of storage time, locust bean gum and dry matter of milk on the physical properties of low-fat set yoghurt. *International Dairy Journal*, 13, 909-916.

Vrhic, N. & Hruskar, M. (2000). Slovenian fermented milk with probiotics. *Zootehnika*, 76, 41-46.

Vasiljevic, T. & Shah, N. P. (2008). Probiotics-From Metchnikoff to bioactives. *International Dairy Journal*, 18, 714-728.

Viljanen, M., Kuitunen, M., Haahtela, T., Juntunen-Backman, K., Korpela, R. & Savilahti, E. (2005). Probiotic effects on faecal inflammatory markers and on faecal IgA in food allergic atopic eczema/dermatitis syndrome infants. *Pediatric Allergy and Immunology*, 16, 65-71.

Vinderola, C. G., and Reinheimer, J. A. (2000). Enumeration of *Lactobacillus casei* in the presence of *Lactobacillus acidophilus*, *bifidobacterium* and lactic starter bacteria in fermented dairy products. *International Dairy Journal*, 10, 271-275.

Vinderola, C. G., Bailo, N. & Reinheimer, J. A. (2000). Survival of probiotic microflora in Argentinian yoghurts during refrigerated storage. *Food Research International*, 33, 97-102.